

Associated charged gauge boson W_H^- and top quark production in the left-right twin higgs model at the LHC

Yao-Bei Liu, Hong-Mei Han, Yong-Hua Cao

Department of Mechanical and Electric Engineering,

Henan Institute of Science and Technology, Xinxiang 453003, P.R.China *

February 2, 2008

Abstract

The twin Higgs mechanism has recently been proposed to solve the little hierarchy problem. In the context of the Left-Right twin Higgs(*LRTH*) model, we discuss single production of the new charged gauge boson W_H^- , which is predicted by the left-right twin Higgs model, in association with top quark at the *CERN* Large Hadron Collider(LHC). It is found that, for a typical nonzero value of mass mixing parameter $M=150\text{GeV}$ in the *LRTH* model, the production cross section is in the range of $3 \times 10^{-2} - 6.07 \times 10^3 \text{fb}$ at the LHC. As long as the W_H^- is not too heavy, the possible signatures of the heavy charged gauge boson might be detected at the LHC experiments.

PACS number(s): 12.60.Cn, 14.70.Hp, 14.80.Bn

*E-mail:hnxxlyb2000@sina.com

1 Introduction

The standard model(SM) provides an excellent effective field theory description of almost all particle physics experiments. But in the SM the Higgs boson mass suffers from an instability under radiative corrections. The naturalness argument suggests that the cutoff scale of the SM is not much above the electroweak scale: New physics will appear around TeV energies. Most of the new physics models predicted the existence of the new gauge bosons with masses in the TeV range. Thus, search for extra gauge bosons provides a common tool in quest for new physics at the next generation collider experiments. The discovery and study of extra gauge bosons is one of the important goals of the *CERN* Large Hadron Collider(LHC).

One interesting approach to the hierarchy problem, first proposed in [1, 2], is that the Higgs mass parameter is protected the pseudo-Goldstone boson of an approximate global symmetry. In the last few years several interesting realizations of this idea based on the little Higgs mechanism have been constructed [3, 4]. These theories stabilize the weak scale up to be above a few TeV. Recently, the twin Higgs mechanism has been proposed as a solution to the little hierarchy problem [5, 6, 7]. The Higgs is a pseudo-Goldstone boson of a spontaneously broken global symmetry. Gauge and Yukawa interactions that explicitly break the global symmetry give mass to the Higgs. Once a discrete symmetry is imposed, the leading quadratic divergent term respects the global symmetry, thus does not contribute to the Higgs model. The resulting Higgs mass obtains logarithmic corrections. Its mass is around the electroweak scale when the cut off is around 5-10 TeV.

The twin Higgs mechanism can be implemented in left-right models with the discrete symmetry being identified with left-right symmetry [6]. The left-right twin Higgs($LRTH$) model contains $U(4)_1 \times U(4)_2$ global symmetry as well as $SU(2)_L \times SU(2)_R \times U(1)_{B-L}$ gauge symmetry. The left-right symmetry acts on only the two $SU(2)$'s gauge symmetry. The leading quadratically divergent SM gauge boson contributions to the Higgs masses are canceled by the loop involving the heavy gauge bosons. A vector top singlet pair is introduced to generate an $\mathcal{O}(1)$ top Yukawa coupling. The quadratically divergent SM top contributions to the Higgs potential are canceled by the contributions from a heavy top partner. The $LRTH$ model predicted the existence of the new heavy particles, such as heavy gauge boson Z_H and $W_H^\pm$, which should not be much heavier than 1 TeV. The characteristic signatures of the $LRTH$ model at

the present and future collider experiments and the production and decay of these new particles have been studied in Ref [8, 9]. It has been shown that the charged gauge bosons from twin Higgs model should be either seen or excluded in the first year of running at the LHC. Thus, it is very interesting to study production of these new charged gauge bosons at the LHC.

In the SM framework, single production of the charged gauge boson $W^\pm$ associated with a top quark is one of the important process for single top quark production. Its cross section is negligible at the Tevatron but of considerable size at the LHC, where it is larger than that of the s-channel process for single top quark production. There is an extensive work on tW associated production at the LHC in the literature [10, 11, 12]. The production cross section for this process has been calculated at leading order, with a subset of the next-to-leading order(NLO) corrections included [10]. Including both the subsequent leptonic decays $W \rightarrow l\nu$ and $t \rightarrow l\nu b$, as well as the emission of real radiation in the top decay, a NLO calculation of this process has been given in Ref.[11]. Recently, Ref.[13] has investigated single production of new gauge bosons associated with a top quark at the LHC in the little Higgs models. In this letter, we will study single production of the charged gauge boson predicted by the $LRTH$ model in association with top quark and see whether the possible signals of this new particle can be detected via this process at the LHC.

This paper is organized as follows, In section two, we first briefly introduce the $LRTH$ model, and then give the production amplitude of the process. The numerical results and discussions are presented in section three. The conclusions are given in section four.

2 The Left-Right twin Higgs model and the production amplitude of $bg \rightarrow tW_H^-$

The $LRTH$ model is based on the global $U(4)_1 \times U(4)_2$ symmetry, with a locally gauged subgroup $SU(2)_L \times SU(2)_R \times U(1)_{B-L}$. The two Higgs fields, H and $\hat{H}$, are introduced and each transforms as $(4, 1)$ and $(1, 4)$ respectively. The introduction of the extra Higgs field $\hat{H}$ enlarge the global symmetry to $U(4)_1 \times U(4)_2$. The global $U(4)_1(U(4)_2)$ symmetry is spontaneously broken down to its subgroup $U(3)_1(U(3)_2)$ with non-zero vacuum expectation values as $\langle H \rangle = (0, 0, 0, f)$ and $\langle \hat{H} \rangle = (0, 0, 0, \hat{f})$. Each spontaneously symmetry breaking results in

seven Nambu-Goldstone bosons. Three of six Goldstone bosons that are charged under $SU(2)_R$ are eaten by the heavy gauge bosons, while leaves three physical Higgs: ϕ^0 and $\phi^\pm$. The remaining Higgses are the SM Higgs doublet H_L and an extra Higgs doublet $\hat{H}_L = (\hat{H}_1^+, \hat{H}_2^0)$ that only couples to the gauge boson sector. A residue matter parity in the model renders the neutral Higgs $\hat{H}_2^0$ stable, and it could be a good dark matter candidate. After electroweak symmetry breaking ($EW\!SB$), a physical Higgs boson h_{SM} from H_L is left, which plays the role of the SM Higgs particle.

The gauge interactions break the global symmetry, which generate a potential for the Goldstone bosons, in particular, for the SM Higgs doublet. The left-right discrete symmetry ensures that the global symmetry is respected at the quadratic order and so the quadratically divergent mass correction contributes only to the masses of the already massive radial modes but not to the masses of the Goldstone bosons. After $EW\!SB$, three of the four $SU(2)_R \times U(1)_{B-L}$ gauge bosons become massive, with masses proportional to $\sqrt{f^2 + \hat{f}^2}$. Since these gauge bosons couple to the SM matter fields, their masses are highly constrained from either precision measurements or direct searches. The large value of $\hat{f}$ does not reintroduce the fine tuning problem for the Higgs potential, since the gauge boson contributions to the Higgs potential is suppressed by the smallness of the gauge couplings.

There are six massive gauge bosons $W^\pm$, $W_H^\pm$, Z , Z_H and one massless photon γ . For the charged gauge bosons, there is no mixing between the $W_L^\pm$ and $W_H^\pm$: $W_L^\pm$ is identified with the SM weak gauge boson $W^\pm$ while $W_H^\pm$ is much heavier than $W^\pm$ and their mass are given to leading order by [8]

$$M_W^2 = \frac{1}{2}g^2 f^2 \sin^2 x, \quad (1)$$

$$M_{W_H}^2 = \frac{1}{2}g^2(\hat{f}^2 + f^2 \cos^2 x), \quad (2)$$

where $x = v/\sqrt{2}f$, the values of f and $\hat{f}$ will be bounded by electroweak precision measurements. In addition, f and $\hat{f}$ are interconnected once we set $v = 246 GeV$. Once f is fixed, the VEV $\hat{f}$ can be determined from the minimization of the CW potential of the SM Higgs. We define the Weinberg angle in the $LRTH$ model as

$$s_w = \sin \theta_w = \frac{g'}{\sqrt{g^2 + 2g'^2}}, \quad (3)$$

$$c_w = \cos \theta_w = \sqrt{\frac{g^2 + g'^2}{g^2 + 2g'^2}}. \quad (4)$$

The unit of the electric charge is then given by

$$e = gs_w = \frac{gg'}{g^2 + 2g'^2}. \quad (5)$$

In order to give the top quark a mass of the order of electroweak scale, a pair of vector-like quarks q_L and q_R are introduced, which are singlets under $SU(2)_L \times SU(2)_R$. The mass of the heavy top is [8]

$$m_T^2 \simeq 2M^2 + m_t^2 \left(\frac{2f^2}{v^2} - 1 \right). \quad (6)$$

Here, the mass parameter M is essential to the mixing between the SM-like top quark and the heavy top quark. The top Yukawa coupling can then be determined by fitting the experimental value of the light top quark mass. At the leading order of $1/f$, the mixing angles for left-handed and right-handed fermions are [8]

$$s_R \simeq \frac{M}{m_T} \left(1 + \frac{v^2}{2f^2} \right). \quad (7)$$

The value of M is constrained by the requirement that the branching ratio of $Z \rightarrow b\bar{b}$ remains consistent with the experiments. It is also constrained by the oblique parameters.

It has been shown that the collinear $\bar{b}$ component of the inclusive process $gg \rightarrow t\bar{b}W^-$ can be attached as a *QCD NLO* correction to bottom quark distribution function of the exclusive process $bg \rightarrow tW^-$ at the partonic level. The *LO* contributions to tW^- associated production is best considered to arise from the process $bg \rightarrow tW^-$. Thus, in this paper, we only consider single production of the heavy charged gauge boson W_H^- in association with a top quark via the two body final state process $bg \rightarrow tW_H^-$ at the LHC as shown in Fig.1. Our numerical results are easily transferred to single production of the heavy gauge boson W_H^+ at the LHC by replacing $\bar{b}$ as b and $\bar{t}$ as t .

At the leading order, the scattering amplitude of the partonic process $bg \rightarrow tW_H^-$ from Fig.1(a) and (b), which can be written as:

$$M = \frac{eg_s s_R}{2\sqrt{2}s_W} \bar{u}(t) \left[\frac{\not{\epsilon}_2(1 + \gamma_5)(\not{P}_g + \not{P}_b + m_b)\not{\epsilon}_1}{\hat{s}' - m_b^2} + \frac{\not{\epsilon}_1(\not{P}_t - \not{P}_g + m_t)\not{\epsilon}_2(1 + \gamma_5)}{\hat{u} - m_t^2} \right] u(b), \quad (8)$$

where $\hat{s}' = (P_g + P_b)^2 = (P_{W_H} + P_t)^2$, $\hat{u} = (P_t - P_g)^2 = (P_b - P_{W_H})^2$. ϵ_1 and ϵ_2 are the gluon and W_H^- polarization vectors, respectively. The constant s_R is the mixing angles for left-handed and right-handed fermions. For a nonzero value of mass mixing parameter M in the *LRT*




Figure 1: Feynman diagrams for the partonic process $bg \rightarrow tW_H^-$

model, its value can be fixed by the symmetry breaking scales f [8].

The cross section for single production of the heavy gauge boson W_H associated with a top quark at the LHC with the center-of-mass $\sqrt{s} = 14TeV$ can be obtained by convoluting the production cross section $\hat{\sigma}(tW_H^-)$ of the partonic process $bg \rightarrow tW_H^-$ with the parton distribution functions ($PDF's$):

$$\sigma(s) = \int_{\tau}^1 dx_1 \int_{\tau/x_1}^1 dx_2 [f_{g/p}(x_1, \mu) f_{b/p}(x_2, \mu) \hat{\sigma}(tW_H^-) + f_{b/p}(x_1, \mu) f_{g/p}(x_2, \mu) \hat{\sigma}(tW_H^-)], \quad (9)$$

where $\tau = (M_{W_H}^2 + m_t^2)/s$ and $\hat{s}' = x_1 x_2 s$. Though out this paper, we will use *CTEQ6L* $PDF's$ [14] for the bottom quark and gluon $PDF's$. Following the suggestions given by Refs.[10, 11, 12], we assume that the factorization scale μ for the bottom quark PDF is of order $(M_{W_H} + m_t)/4$.

3 The numerical results and discussions

In the numerical calculation, we take the input parameters as $m_t = 171.4$ GeV [15], $\alpha_e = 1/128.8$, $\alpha_s = 0.118$ and $s_W^2 = 0.2315$ [16]. Except for these *SM* input parameters, the production cross sections of the new charged gauge boson from twin Higgs model is dependent on the symmetry breaking scale f and the mass of the heavy gauge boson W_H . Taking into account the precision electroweak constraints on the parameter space, the symmetry breaking scales f is allowed in the range of $f = 500 \sim 1500GeV$. While the mass of the heavy gauge boson W_H is larger than 1 TeV [8]. As numerical estimation, we will take M_{W_H} as a free parameter and assume it in the range of $1 \sim 4TeV$.




Figure 2: The cross section $\sigma(s)$ as a function of the mass parameter M_{W_H} for $M=150$ GeV and different values of the symmetry breaking scales $f=500, 1000$ and 1500 GeV.

In Fig.2, we plot the cross section $\sigma(s)$ for single production of the new gauge boson W_H^- associated with top quark at the LHC as a function of the mass parameter M_{W_H} for $M=150$ GeV and three values of the symmetry breaking scales f . One can see from Fig.2 that the cross section $\sigma(s)$ increase as the symmetry breaking scales f decrease. For $f=1000$ GeV and $1\text{TeV} \leq M_{W_H} \leq 3\text{TeV}$, the value of the cross section $\sigma(s)$ is in the range of $1.3 \times 10^3 \sim 1.2\text{fb}$. If we assume the yearly integrated luminosity of the LHC is $\mathcal{L} = 100\text{fb}^{-1}$, then there could be as much as hundreds of events produced each year.

To see the influence of the symmetry breaking scale f on the cross section, in Fig.3 we plot $\sigma(s)$ as a function of the symmetry breaking scale f for $M=150$ GeV and three values of $M_{W_H} = 2, 2.5, 3\text{TeV}$, respectively. From Fig.3, one can see that the cross section decreases as f increases for fixed new gauge boson mass M_{W_H} . This is because the value of the production section is proportional to the order of $1/f^2$, which comes from the flavor change couplings $g^{W_H tb}$. On the other hand, as long as the heavy gauge boson mass M_{W_H} is smaller than 2 TeV, the cross section can even reach hundreds of fb. If we take integral luminosity $\mathcal{L} = 100\text{fb}^{-1}$, there are about $10^3 - 10^5$ events to be produced. There will be a promising number of fully reconstructible events to detect at the LHC.

The cross section depends sensitively on the parameter M , which is the mass mixing




Figure 3: The cross section $\sigma(s)$ of single W_H production as a function of the symmetry breaking scales f for $M=150$ GeV and different values of M_{W_H} .




Figure 4: The cross section $\sigma(s)$ of single W_H production as a function of the mass parameter M_{W_H} for $f=1$ TeV and three different values of the parameter M .

between the SM-like top quark and the heavy top quark. In Fig.4, we plot the cross sections as the function of the mass parameter M_{W_H} for three values of the parameter M , and take $f=1$

TeV as the examples. From Fig.4, one can see that the cross section decreases as the parameter M decreases for fixed the symmetry breaking scale f . Our results do not change much if other small values of M is used. However, once M is very small or in the limit that $M=0$, certain couplings go to zero, which changes the collider signatures significantly. Furthermore, it yields large log divergence of the SM Higgs mass. To compensate for it the heavy gauge bosons also get large masses, it is natural for us to take a typical nonzero value of the mass mixing parameter $M=150\text{GeV}$ in Fig.2 and Fig.3.

To see whether the heavy gauge boson W_H^- can be observed at the LHC via the process $pp \rightarrow gb + X \rightarrow tW_H^- + X$, we discuss the possible decay modes of the heavy gauge boson. For W_H^- , the dominant decay into two jets is not very useful due to the huge QCD jet background. W_H could decay into $l\nu_R$ if the right handed neutrino mass is less than M_{W_H} . The reconstruction of ν_R can be used to reconstruct the W_H mass and distinguish the signal from the background. It leads to a clean signal of lepton plus missing energy. W_H could also decay into a heavy top plus a b -jet, with a branching of about $20\% - 30\%$ [8]. Depending on the subsequent decays of the heavy top, we expect to see signals of $4b + \text{lepton}(e \text{ or } \mu) + \text{missing } E_T$ or $2b + \text{tri-lepton} + \text{missing } E_T$. Although the signal is smaller due to the small leptonic Z, W decay branching ratios and small $Br(T \rightarrow Zt)$, the tri-lepton signal provides a very nice suppression of the SM background. The $W_H \rightarrow tb$ branching ratio is of the order of 4% or less, which has been studied in Ref.[17]. It has been shown that at the LHC, with $10 - 100\text{fb}^{-1}$ luminosity, a reach of $3 - 4$ TeV is possible at 95% C.L.

4 Conclusion

The twin Higgs mechanism provides an alternative method to solve the little hierarchy problem. The $LRTH$ model is a concrete realization of the twin Higgs mechanism. The cancelations of divergences occurs by alignment of vacua and existence of several new particles. The new particles in the $LRTH$ model are heavy top quark, new gauge bosons, and several scalars. In this letter, we investigate single production of new charged gauge boson W_H^- , which is predicted by the $LRTH$ model, can be abundantly produced at the LHC. The collider phenomenology of the $LRTH$ model depends sensitively on the parameter M , which is the mass mixing between

the SM-like top quark and the heavy top quark. For a typical nonzero value of the mass mixing parameter $M=150\text{GeV}$, our results show that as long as the new charged gauge boson is not too heavy, the possible signatures of the heavy charged gauge boson might be detected at the LHC experiments. Certainly, more detailed analysis of the SM background would be needed to make a quantitative conclusion for the W_H^- observation.

References

- [1] H. Georgi and A. Pais, *Phys. Rev.* D**10** (1974) 539; *Phys. Rev.* D**12** (1975) 508.
- [2] D. B. Kaplan and H. Georgi, *Phys. Lett.* B**136** (1984) 183; D. B. Kaplan, H. Georgi and S. Dimopoulos, *Phys. Lett.* B**136** (1984) 187; H. Georgi and D. B. Kaplan, *Phys. Lett.* B**145** (1984) 216.
- [3] N. Arkani-Hamed, A. G. Cohen, and H. Georgi, *Phys. Lett.* B**513** (2001) 232.
- [4] N. Arkani-Hamed, A. G. Cohen, E. Katz, A. E. Nelson, T. Gregoire, and J. G. Wacker, *JHEP* **0208** (2002) 021; I. Low, W. Skiba, and D. Smith, *Phys. Rev.* D**66** (2002) 072001; D. E. Kaplan and M. Schmaltz, *JHEP* **0310** (2003) 039.
- [5] Z. Chacko, H. S. Goh and R. Harnik, *Phys. Rev. Lett* **96** (2006) 231802; Z. Chacko, Y. Nomura, M. Papucci and G. Perez, *JHEP* **0601** (2006) 126.
- [6] Z. Chacko, H. S. Goh and R. Harnik, *JHEP* **0601**(2006) 108.
- [7] A. falkowski, S. Pokorski and M. Schmaltz, *Phys. Rev.* D**74** (2006) 035003.
- [8] Hock-Seng Goh, Shufang Su, hep-ph/0611015.
- [9] Dong-Won Jung , Jae-Yong Lee, hep-ph/0701071; Yao-Bei Liu, Hong-Mei Han and Xue-Lei Wang, hep-ph/0701169.
- [10] A. P. Heinson, A. S. Belyaev, E. E. Boos, *Phys. Rev.* D**56** (1997) 3114; T. M. P. Tait, *Phys. Rev.* D**61** (2000) 034001; A. Belyaev, E. Boos, *Phys. Rev.* D**63** (2001) 034012.
- [11] J. Campbell and F. Tramontano, *Nucl. Phys.* B**726** (2005) 109.
- [12] M. Beccaria, F. M. Denard, C. Verzegnassi, *Phys. Rev.* D**71** (2005) 033005; M. Beccaria et al, *Phys. Rev.* D**73** (2006) 093001.
- [13] Chong-Xing Yue, Shuo Yang and Li-Hong Wang, *Europhys. Lett.* **76** (2006) 381-387.
- [14] J. Pumplin et al., *JHEP* **0207** (2002) 012; D. Stimp et al., *JHEP* **0310** (2003) 046.

- [15] The Tevatron Electroweak Working Group for CDF and D0 Collaborations, *Combination of CDF and D0 Results on the Mass of the Top Quark*, hep-ex/0608032.
- [16] Particle Data Group, D. E. Groom *et al.* *Eur.Phys.J.C* **15** (2000) 1.
Particle Data Group, K. Hagiwara *et al.*, *Phys. Rev.* **D66** (2002) 010001.
- [17] Z. Sullivan, *Phys. Rev.* **D66** (2002) 075011.